

**DIRECTIONS FOR GASTROGRAFIN  
(USED FOR CT IN PLACE OF BARIUM)**

NAME: \_\_\_\_\_

**APPOINTMENT INFORMATION**

DATE: \_\_\_\_\_ TIME: \_\_\_\_\_

TAKE 2 SYRINGES FULL OF THE CONTRAST. EACH ONE MIXED WITH EITHER 8 OZ. OF APPLE OR WHITE GRAPE JUICE.

DRINK 1<sup>ST</sup> SYRINGE AND JUICE MIX **2 HOURS** PRIOR TO THE EXAM

DRINK 2<sup>ND</sup> SYRINGE AND JUICE MIX **½ HOUR** PRIOR TO YOUR EXAM.

**NO SOLID FOOD 3 HOURS PRIOR TO THE EXAM.**

---

**DIRECTIONS FOR GASTROGRAFIN  
(USED FOR CT IN PLACE OF BARIUM)**

NAME: \_\_\_\_\_

**APPOINTMENT INFORMATION**

DATE: \_\_\_\_\_ TIME: \_\_\_\_\_

TAKE 2 SYRINGES FULL OF THE CONTRAST. EACH ONE MIXED WITH EITHER 8 OZ. OF APPLE OR WHITE GRAPE JUICE.

DRINK 1<sup>ST</sup> SYRINGE AND JUICE MIX **2 HOURS** PRIOR TO THE EXAM

DRINK 2<sup>ND</sup> SYRINGE AND JUICE MIX **½ HOUR** PRIOR TO YOUR EXAM.

**NO SOLID FOOD 3 HOURS PRIOR TO THE EXAM.**

---

**DIRECTIONS FOR GASTROGRAFIN  
(USED FOR CT IN PLACE OF BARIUM)**

NAME: \_\_\_\_\_

**APPOINTMENT INFORMATION**

DATE: \_\_\_\_\_ TIME: \_\_\_\_\_

TAKE 2 SYRINGES FULL OF THE CONTRAST. EACH ONE MIXED WITH EITHER 8 OZ. OF APPLE OR WHITE GRAPE JUICE.

DRINK 1<sup>ST</sup> SYRINGE AND JUICE MIX **2 HOURS** PRIOR TO THE EXAM

DRINK 2<sup>ND</sup> SYRINGE AND JUICE MIX **½ HOUR** PRIOR TO YOUR EXAM.

**NO SOLID FOOD 3 HOURS PRIOR TO THE EXAM.**